Cypress Swamp



Status

Current Condition: Poor and declining. According to the best available GIS information at this time (see Appendix C: GIS Data Tables), 1,586,941 acres (642,212 ha) of Cypress Swamp habitat exist, of which 44% (689,955 ac; 279,215 ha) are in existing conservation or managed areas. Another 11% (173,971 ac; 70,404 ha) are in Florida Forever projects and 10% (163,702 ac; 66,248 ha) are in SHCA-designated lands. The remaining 35% (559,313 ac; 226,346 ha) are other private lands.



Some habitat distributions or locations may be misrepresented on this map due to size, resolution and insufficient data sources.

Habitat Description

FNAI type: Strand Swamp, Dome Swamp

These regularly inundated wetlands form a forested border along large rivers, creeks, and lakes, or occur in depressions as circular domes or linear strands. These communities are strongly dominated by either bald cypress or pond cypress, with very low numbers of scattered black gum, red maple, and sweetbay. Understory and ground cover are usually sparse due to frequent flooding but sometimes include such species as buttonbush, lizard's-tail, and various ferns.

Associated Species of Greatest Conservation Need

Mammals

- Corynorhinus rafinesquii
- Eumops floridanus

Rafinesque's Big-eared Bat Florida Bonneted Bat Lasiurus borealis borealis

Lasiurus intermedius floridanus

Lasiurus seminolus

Myotis austroriparius

Perimyotis subflavus

Lontra canadensis lataxina

Neovison vison evergladensis

Neovison vison halilimnetes

Puma concolor coryi

Ursus americanus floridanus

Trichechus manatus latirostris

Red Bat

Northern Yellow Bat

Seminole Bat

Southeastern Myotis

Tricolored Bat

River Otter

Everglades Mink

Gulf Salt Marsh Mink

Florida Panther

Florida Black Bear

West Indian Manatee

Birds

Mycteria americana

Ardea herodias

Ardea alba

Egretta thula

Egretta caerulea

Butorides virescens

Nycticorax nycticorax

Nyctanassa violacea

Eudocimus albus

Plegadis falcinellus

Elanoides forficatus

Haliaeetus leucocephalus

Buteo brachyurus

Aramus guarauna

Campephilus principalis

Vermivora chrysoptera

Vermivora cyanoptera

Protonotaria citrea

Setophaga ruticilla

Setophaga dominica stoddardi

Setophaga discolor discolor

Cardellina canadensis

Euphagus carolinus

Wood Stork

Great Blue Heron

Great Egret

Snowy Egret Little Blue Heron

Green Heron

Black-crowned Night-Heron

Yellow-crowned Night-Heron

White Ibis

Glossy Ibis

Swallow-tailed Kite

Bald Eagle

Short-tailed Hawk

Limpkin

Ivory-billed Woodpecker

Golden-winged Warbler

Blue-winged Warbler

Prothonotary Warbler

American Redstart

Stoddard's Yellow-throated Warbler

Reticulated Flatwoods Salamander

Frosted Flatwoods Salamander

Prairie Warbler

Canada Warbler

Rusty Blackbird

Gopher Frog Carpenter Frog

Ornate Chorus Frog

Amphibians

Lithobates capito

Lithobates virgatipes

Pseudacris ornata

Ambystoma bishopi

Ambystoma cingulatum

Ambystoma tigrinum

Amphiuma pholeter

Desmognathus auriculatus

Eurycea chamberlaini Hemidactylium scutatum

Notophthalmus perstriatus

Pseudobranchus striatus lustricolus

Pseudobranchus striatus striatus Stereochilus marginatus

Southern Dusky Salamander

Chamberlain's Dwarf Salamander

Eastern Tiger Salamander

One-toed Amphiuma

Four-toed Salamander

Striped Newt

Gulf Hammock Dwarf Siren

Broad-striped Dwarf Siren

Many-lined Salamander

Reptiles

Alligator mississippiensis
Anolis carolinensis seminolus
Plestiodon anthracinus pluvialis
Crotalus horridus
Drymarchon couperi
Farancia erytrogramma
American Alligator
Southern Green Anole
Southern Coal Skink
Timber Rattlesnake
Eastern Indigo Snake
Rainbow Snake

Heterodon platirhinos
Lampropeltis getula
Eastern Hog-nosed Snake
Eastern Kingsnake

Nerodia cyclopion
Seminatrix pygaea cyclas
Mississippi Green Watersnake
Southern Florida Swampsnake

Clemmys guttata Spotted Turtle
Deirochelys reticularia Chicken Turtle
Terrapene carolina Eastern Box Turtle

Fish

Hybognathus hayi
Notropis melanostomus
Pteronotropis welaka
Umbra pygmaea
Atractosteus spatula
Acantharchus pomotis
Enneacanthus chaetodon
Cypress Minnow
Black mouth Shiner
Eastern Mudminnow
Alligator Gar
Mud Sunfish
Black Banded Sunfish

Enneacanthus chaetodon
Etheostoma proeliare
Cypress Darter

Invertebrates

Cambarellus blacki Cypress Crayfish A Crayfish Cambarellus schmitti A Crayfish Procambarus apalachicolae A Crayfish Procambarus latipleurum Tail-light Damsel Chrysobasis lucifer Lestes tenuatus Blue-striped Spreadwing Euphyes berryi Berry's Skipper Euphyes dion Dion Skipper Hesperia attalus slossonae Seminole Skipper Callophrys henrici Henry's Elfin Callophrys hesseli Hessel's Hairstreak Zale perculta Okefenokee Zale Moth Seminole Crescent Anthanassa texana seminole Enodia portlandia floralae Florida Pearly Eye

Conservation Threats

Threats to the Cypress Swamp habitat that were also identified for multiple other habitats are addressed in Chapter 7: Multiple Habitat Threats and Conservation Actions. These threats include:

- Conversion to agriculture
- Conversion to housing and urban development
- Groundwater withdrawal
- Incompatible fire
- Incompatible forestry practices

- Incompatible resource extraction—mining/drilling
- Invasive animals
- Invasive plants
- Nutrient loads-agriculture

- Nutrient loads—urban
- Roads
- Surface water withdrawal and diversion

Widespread ditching and diking of this habitat and hydrologic fragmentation due to construction of roads through and adjacent to this habitat are large sources of altered hydrologic regime. Groundwater withdrawal for municipal and agricultural purposes has impacted cypress wetlands in localized areas throughout Florida, but this threat is most severe in portions of central Florida. Incompatible forestry practices threaten this habitat due to physical and hydrological disturbance and the slow regeneration time of cypress trees. Currently, most cypress harvest is of young, small-diameter trees for landscape mulch. Nearly all cypress wetlands in unprotected lands have suffered from altered landscape context as the surrounding uplands and wet prairies have been converted to other land uses, primarily agriculture and urban/suburban development. In many parts of Florida, cypress wetlands are particularly vulnerable to and have been seriously impacted by a variety of invasive plants. Many cypress wetlands in both agricultural and urban settings receive nutrient-laden discharges from stormwater management systems, often leading to drastic changes in understory plant community composition and associated faunal changes. Additional threats specific to this habitat include the numerous water control structures affecting Cypress Swamps, particularly smaller dome swamps, statewide.

The following stresses and sources of stress threaten this habitat:

	Stresses	Habitat Stress Rank
A	Altered hydrologic regime	High
В	Altered landscape mosaic or context	High
C	Altered soil structure and chemistry	High
D	Altered community structure	High
Е	Altered species composition/dominance	High
F	Habitat destruction or conversion	Medium
G	Altered water quality of surface water or aquifer: nutrients	Medium
Н	Missing key communities, functional guilds, or seral stages	Medium
I	Altered fire regime	Medium
J	Fragmentation of habitats, communities, ecosystems	Medium
K	Altered water and/or soil temperature	Low
L	Habitat degradation/disturbance	Low

The sources of stress, or threats, were used to generate conservation actions.

Sources of Stress		Habitat Source Rank	Related Stresses (see above)
1	Incompatible forestry practices	High	A, B, C, D, E, F, H
2	Surface water withdrawal	High	A, B, C, D, E, F
3	Nutrient loads-agriculture	High	E, G
4	Invasive plants	High	D, E
5	Conversion to housing and urban development	High	A, B
6	Invasive animals	Medium	C, D, E
7	Groundwater withdrawal	Medium	A, C, E
8	Roads	Medium	A, B, E
9	Conversion to agriculture	Medium	A, B
10	Incompatible vegetation harvest	Low	Е
11	Nutrient loads-urban	Low	E, G
12	Incompatible fire	Low	B, E
13	Incompatible resource extraction: mining/drilling	Low	A, F
14	Incompatible grazing and ranching	Low	D, E, G
15	Incompatible agricultural practices	Low	A
16	Management of nature–water control structures	Low	A, B
State	Statewide Threat Rank of Habitat		

Conservation Actions

Actions to abate the threats to Cypress Swamp that were also identified as statewide threats (incompatible forestry practices, surface water withdrawal and diversion, nutrient loads—agriculture, invasive plants, conversion to housing and urban development, invasive animals, groundwater withdrawal, roads, conversion to agriculture, nutrient loads—urban, incompatible fire, and incompatible resource extraction—mining/drilling) are in Chapter 7: Multiple Habitat Threats and Conservation Actions.

Several of the actions developed for a statewide threat that were only applicable to Cypress Swamp and a few other habitats (i.e., Aquatic Cave, Calcareous Stream, Freshwater Marsh and Wet Prairie, Natural Lake, Reservoir/Managed Lake, Seepage/Steephead Stream, Softwater Stream, Spring and Spring Run, Terrestrial Cave, and Coastal Tidal River or Stream) and are listed below. Additional actions were developed to address threats specific to this habitat. These actions are intended to increase the spatial extent of Cypress Swamps in the landscape and improve the functionality of existing cypress wetlands through both regional and small-scale hydrologic restoration projects.

Incompatible Forestry Practices

	J.			
Overall Rank	Education and Awareness	Feasibility	Benefits	Cost
L	Encourage labeling on cypress mulch alternatives that promotes their ecological value to consumers.	M	L	L
L	Through garden clubs, landscapers, and other avenues, promote acceptable alternatives to cypress mulch and make them readily available.	M	L	M
Overall Rank	Research	Feasibility	Benefits	Cost
L	Investigate various sources of possible funding for cypress regeneration studies	M	L	L
L	Recognizing that species move between wetland and upland habitats, assess the effectiveness of current BMP's regarding bedding near isolated wetlands.	M	L	L

Conversion to Housing and Urban Development

Overall Rank	Economic and Other Incentives	Feasibility	Benefits	Cost
L	Encourage tax or other incentives, such as density transfers, for environmentally friendly comprehensive development plans for projects that front on rivers and floodplains.	M	L	VH

Conversion to Agriculture

Overall Rank	Economic and Other Incentives	Feasibility	Benefits	Cost
M	Create incentives for maintenance and conversion of lands to agricultural uses that use less water and result in lower nutrient outputs into Florida's waters and wetlands, and create market-based incentives to compensate private landowners for the environmental services they provide to the state through management that increases water storage and nutrient reduction.	M	M	Н

Management of Nature – Water Control Structures

Overall Rank	Economic and Other Incentives	Feasibility	Benefits	Cost
M	Review existing Farm Bill programs and explore options for enhancing economic benefits to landowners that improve or remove water control structures.	VH	L	L
Overall Rank	Education and Awareness	Feasibility	Benefits	Cost
L	Develop an awareness program for Drainage Districts created by Chapter 298 of the Florida Administrative Code ("298 Districts") to educate them about opportunities to improve fish and wildlife habitat conditions through operational and/or structural changes in their drainage systems.	Н	L	M
Overall Rank	Land/Water/Species Management	Feasibility	Benefits	Cost
L	Create a grant program (or utilize existing Farm Bill and other federal programs) to replace or retrofit existing stop log or manually controlled structures with V-notch weirs in agricultural drainage systems. Give priority to those control structures that are identified as acting as barriers to wildlife movement or sheet flow.	Н	L	Н
Overall Rank	Policy	Feasibility	Benefits	Cost